

Immunization Schedule with Combination Vaccines

EVERY FALL: FLU VACCINE⁴ for anyone 6 months and older

	2 MONTHS	4 MONTHS	6 MONTHS	12 MONTHS	15 MONTHS	18 MONTHS	4-6 YEARS		
Schedule Using PEDIARIX[®] PROQUAD[®] KINRIX[™]	PEDIARIX[®] DTaP, IPV, HepB + PCV Rotavirus Hib	PEDIARIX[®] DTaP, IPV, HepB ¹ + PCV Rotavirus Hib	PEDIARIX[®] DTaP, IPV, HepB + PCV Rotavirus ² Hib ³	PCV Hib HepA MMR ⁶ Varicella ⁶	DTaP	HepA	KINRIX^{®5} DTaP, IPV + PROQUAD[®] MMRV		
	Schedule Using PENTACEL^{®5} PROQUAD[®] KINRIX[™]	PENTACEL[®] DTaP, IPV, Hib + PCV Rotavirus HepB	PENTACEL[®] DTaP, IPV, Hib + PCV Rotavirus HepB ¹	PENTACEL[®] DTaP, IPV, Hib + PCV Rotavirus ² HepB	PCV HepA MMR ⁶ Varicella ⁶	PENTACEL[®] DTaP, IPV, Hib	HepA	KINRIX^{®5} DTaP, IPV + PROQUAD[®] MMRV	
		Schedule Using PROQUAD[®] & KINRIX[™]	DTaP IPV HepB Hib PCV Rotavirus	DTaP IPV HepB ¹ Hib PCV Rotavirus	DTaP IPV HepB Hib ³ PCV Rotavirus ²	Hib PCV HepA MMR ⁶ Varicella ⁶	DTaP	HepA	KINRIX^{™5} DTaP, IPV + PROQUAD[®] MMRV

Make sure the vaccine you administer contains the antigens on the doctor's order. Keep it simple. Stick with the same product.

This is a suggested schedule for VFC providers ordering combination vaccines. For alternatives and details, consult the latest "Recommended Immunization Schedules for persons aged 0-18 years, United States." For more info, visit eziz.org

¹ A dose of Hepatitis B vaccine is not necessary at 4 months if doses are given at birth and 2 months but may be included as part of a combination vaccine.
² The six month dose is not needed if Rotarix[®] was used exclusively for both dose 1 and 2 of the rotavirus vaccine series.
³ This six month Hib dose is not indicated if PedvaxHIB[®] is used exclusively for the 2 and 4 month infant doses.
⁴ Influenza vaccine is available in thimerosal-free options. See California Health and Safety Code § 124172.

⁵ Licensed by FDA for children 4 through 6 years with previous doses of INFANRIX[™] or PEDIARIX[™]. ACIP recommends that, whenever feasible, the same manufacturer's DTaP vaccines be used for each dose in the series; however, vaccination should not be deferred because the type of DTaP previously administered is unavailable or unknown. See www.cdc.gov/mmwr/preview/mmwrhtml/mm5739a4.htm
⁶ CDC recommends MMR + Varicella at 12-15 months. Providers can use their discretion whether to use MMRV, however.

